No.



200300238

<u>THE UNITED STATES OF ANTERIOR</u>

TO ALL TO WHOM THESE PRESENTS SHALL COME;

Hioneer Hi-Bred International, Inc.

MICCORS, THERE HAS BEEN PRESENTED TO THE

Secretary of Agriculture

AN APPLICATION REQUESTING A CERTIFICATE OF PROTECTION FOR AN ALLEGED DISTINCT VARIETY OF SEXUALLY REPRODUCED, OR TUBER PROPAGATED PLANT, THE NAME AND DESCRIPTION OF WHICH ARE CONTAINED IN THE APPLICATION AND EXHIBITS, A COPY OF WHICH IS HEREUNTO ANNEXED AND MADE A PART HEREOF, AND THE various requirements of LAW in such cases made and provided have been complied with, and the TITLE THERETO IS, FROM THE RECORDS OF THE PLANT VARIETY PROTECTION OFFICE, IN THE APPLICANT(S) INDICATED IN THE SAID COPY, AND WHEREAS, UPON DUE EXAMINATION MADE, THE SAID APPLICANT(S) IS (ARE) ADJUDGED TO BE ENTITLED TO A CERTIFICATE OF PLANT VARIETY PROTECTION UNDER THE LAW.

NOW, THEREFORE, THIS CERTIFICATE OF PLANT VARIETY PROTECTION IS TO GRANT UNTO THE SAID APPLICANT(S) AND THE SUCCESSORS, HEIRS OR ASSIGNS OF THE SAID APPLICANT(S) FOR THE TERM OF TWENTY YEARS FROM THE DATE OF THIS GRANT, SUBJECT TO THE PAYMENT OF THE REQUIRED FEES AND PERIODIC REPLENISHMENT OF VIABLE BASIC SEED OF THE VARIETY IN A PUBLIC REPOSITORY AS PROVIDED BY LAW, THE RIGHT TO EXCLUDE OTHERS OM SELLING THE VARIETY, OR OFFERING IT FOR SALE, OR REPRODUCING IT, OR IMPORTING IT, OR EXPORTING IT, INDITIONING IT FOR PROPAGATION, OR STOCKING IT FOR ANY OF THE ABOVE PURPOSE , OR CONDITIONING IT PAGATION, OR STOCKING IT FOR ANY OF THE ABOVE PURPOSE, OR USING IT IN PRODUCING A HYBRID OR VARIETY THEREFROM, TO THE EXTENT PROVIDED BY THE PLANT VARIETY PROTECTION ACT. (84

CORN, FIELD

'PHR18'

In Jestimonn Thereof, I have hereunto set my hand and caused the seal of the Hant Bariety Frotection Office to be affixed at the City of Washington, D.C. this nineteenth day of August,

in the year two thousand and four.

AMENDED, 7 U.S.C. 2321 ET SEQ.)

(See reverse for instructions and infor-

T-470 (02-10-2003) designed by the Plant Variety Protection Office using Word 2000. Replaces former versions of ST-470, which are obsolete

SENERAL: To be effectively filed with the Plant Variety Protection Office (PVPO), ALL of the following items must be received in the PVPO: (1) Completed application form signed by the owner; (2) completed exhibits A, B, C, E; (3) for a seed reproduced variety at least 2,500 viable untreated seeds, for a hybrid rariety at least 2,500 untreated seeds of each line necessary to reproduce the variety, or for tuber reproduced varieties verification that a viable (in the sense that t will reproduce an entire plant) tissue culture will be deposited and maintained in an approved public repository; (4) check drawn on a U.S. bank for \$3,652 (\$432 iling fee and \$3,220 examination fee), payable to "Treasurer of the United States" (See Section 97.6 of the Regulations and Rules of Practice.) Partial applications will be held in the PVPO for not more than 90 days, then returned to the applicant as unfiled. Mail application and other requirements to Plant Variety Protection Office, AMS, USDA, Room 401, NAL Building, 10301 Baltimore Avenue, Beltsville, MD 20705-2351. Retain one copy for your files. All items on the ace of the application are self explanatory unless noted below. Corrections on the application form and exhibits must be initialed and dated. DO NOT use nasking materials to make corrections. If a certificate is allowed, you will be requested to send a check payable to "Treasurer of the United States" in the amount of \$432 for issuance of the certificate. Certificates will be issued to owner, not licensee or agent.

> **Plant Variety Protection Office** Telephone: (301) 504-5518 FAX: (301) 504-5291

Homepage: http://www.ams.usda.gov/science/pvpo/pvp.htm

TEM

- 18a. Give: (1) the genealogy, including public and commercial varieties, lines, or clones used, and the breeding method:
 - (2) the details of subsequent stages of selection and multiplication;
 - (3) evidence of uniformity and stability; and
 - (4) the type and frequency of variants during reproduction and multiplication and state how these variants may be identified
- 18b. Give a summary of the variety's distinctness. Clearly state how this application variety may be distinguished from all other varieties in the same crop. If the new variety is most similar to one variety or a group of related varieties:
 - (1) identify these varieties and state all differences objectively;
 - (2) attach statistical data for characters expressed numerically and demonstrate that these are clear differences; and
 - (3) submit, if helpful, seed and plant specimens or photographs (prints) of seed and plant comparisons which clearly indicate distinctness.
- 18c. Exhibit C forms are available from the PVPO Office for most crops; specify crop kind. Fill in Exhibit C (Objective Description of Variety) form as completely as possible to describe your variety.
- 18d. Optional additional characteristics and/or photographs. Describe any additional characteristics that cannot be accurately conveyed in Exhibit C. Use comparative varieties as is necessary to reveal more accurately the characteristics that are difficult to describe, such as plant habit, plant color, disease resistance, etc.
- 18e. Section 52(5) of the Act requires applicants to furnish a statement of the basis of the applicant's ownership. An Exhibit E form is available from the PVPO.
- If "Yes" is specified (seed of this variety be sold by variety name only, as a class of certified seed), the applicant MAY NOT reverse this affirmative decision after the variety has been sold and so labeled, the decision published, or the certificate issued. However, if "No" has been specified, the applicant may change the choice. (See Regulations and Rules of Practice, Section 97.103).
- See Sections 41, 42, and 43 of the Act and Section 97.5 of the regulations for eligibility requirements.
- 23. See Section 55 of the Act for instructions on claiming the benefit of an earlier filing date.
- CONTINUED FROM FRONT (Please provide a statement as to the limitation and sequence of generations that may be certified.)
- CONTINUED FROM FRONT (Please provide the date of first sale, disposition, transfer, or use for each country and the circumstances, if the variety including any harvested material) or a hybrid produced from this variety has been sold, disposed of, transferred, or used in the U.S. or other countries.)

United States, Turkey, Nov. 1, 2002.

CONTINUED FROM FRONT (Please give the country, date of filing or issuance, and assigned reference number, if the variety or any component of the variety is protected by intellectual property right (Plant Breeder's Right or Patent).)

It is the responsibility of the applicant/owner to keep the PVPO informed of any changes of address or change of ownership or assignment or owner's epresentative during the life of the application/certificate. There is no charge for filling a change of address. The fee for filling a change of ownership or assignment or any modification of owner's name is specified in Section 97.175 of the regulations. (See Section 101 of the Act, and Sections 97.130, 97.131, 37.175(h) of the Regulations and Rules of Practice.)

To avoid conflict with other variety names in use, the applicant must check the appropriate recognized authority. For example, for agricultural and regetable crops, contact: Seed Branch, AMS, USDA, Room 213, Building 306, Beltsville Agricultural Research Center--East, Beltsville, MD 20705. [elephone: (301) 504-8089. http://www.ams.usda.gov/lsg/seed.htm

According to the Paperwork Reduction Act of 1995, an agency may not conduct or sponsor, and a person is not required to respond to a collection of information unless it displays a valid OMB control number. The alid OMB control number for this information collection is 0581-0055. The time required to complete this information collection is estimated to average 3.0 hours per response, including the time for reviewing nstructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information.

he U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, gender, religion, age, disability, sexual orientation, marital or family tatus, political beliefs, parental status, or protected genetic information. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program nformation (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at 202-720-2600 (voice and TDD).

o file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326-W, Whitten Building, 14th and Independence Avenue, SW, Washington, DC 20250-9410 or call 202-720-5964 (voice and

DD). USDA is an equal opportunity provider and employer.

T-470 (02-10-2003) designed by the Plant Variety Protection Office with Word 2000. Replaces former versions of STS-470, which are obsolete.

Exhibit A. Origin and Breeding History

Pedigree: PH1B5/PH2BB)CX42211

Pioneer Line PHB18, Zea mays L., a flint-dent corn inbred, was developed by Pioneer Hi-Bred International, Inc. from the single cross hybrid PH1B5 (Certificate No. 9800355) X PH2BB using the pedigree method of plant breeding. Varieties PH1B5 and PH2BB are proprietary inbred lines of Pioneer Hi-Bred International, Inc. Variety PH2BB was derived by pedigree selection from the single cross hybrid PHR03 (PVP Certificate No. 9100097) X PHBE2 (PVP certificate No. 9500200).

Selfing was practiced from the above hybrid for 7 generations using pedigree selection. During line development, crosses were made to inbred testers for the purpose of estimating the line's combining ability. Yield trials were grown at Garden City, Kansas as well as other Pioneer research locations. After initial testing, additional hybrid combinations have been evaluated and subsequent generations of the line have been grown and hand-pollinated with observations again made for uniformity.

Variety PHB18 has shown uniformity and stability for all traits as described in Exhibit C - "Objective Description of Variety". It has been self-pollinated and ear-rowed 5 generations with careful attention paid to selection criteria and uniformity of plant type to assure genetic homozygousity and phenotypic stability. The line has been increased both by hand and in isolated fields with continued observations for uniformity and stability, and for 3 generations during the final stages of inbred development and seed multiplication. Very high standards for genetic purity have been established morphologically using field observations and electrophoretically using sound lab molecular marker methodology.

No variant traits have been observed or are expected in PHB18.

The criteria used in the selection of PHB18 were yield, both per se and in hybrid combinations; late season plant health, grain quality, stalk lodging resistance, and kernel size, especially important in production. Other selection criteria include: ability to germinate in adverse conditions; disease and insect resistance; pollen yield and tassel size.

Exhibit A: Developmental history for PHB18

Season/Year Pedigree Grown	Inbreeding Level of Pedigree Grown
PH1B5	F0
PH2BB	F0
PH1B5/PH2BB WINTER 1996	F1
PH1B5/PH2BB)CX SUMMER 1997	F2
PH1B5/PH2BB)CX4 SUMMER 1998	F3
PH1B5/PH2BB)CX41 WINTER 1998	F4
PH1B5/PH2BB)CX412 SUMMER 1999	F5
PH1B5/PH2BB)CX4121 WINTER 1999	F6
PH1B5/PH2BB)CX41211 SUMMER 2000	F7
PH1B5/PH2BB)CX41211X	F8

^{*}PHB18 was selfed and ear-rowed from F3 through F7 generation. #Uniformity and stability were established from F5 through F7 generation and beyond when seed supplies were increased.

Exhibit B: Novelty Statement

Variety PHB18 mostly resembles Pioneer Hi-Bred International, Inc. proprietary inbred line PHBE2 (PVP Certificate No. 9600201). Tables 1A and 1B show two sample t-tests on data collected primarily in Johnston, and Dallas Center, IA. The traits collectively show measurable differences between the two varieties.

Exhibit B: Novelty Statement

Variety PHB18 has a greater husk length (23.5 cm vs 19.3 cm) than variety PHBE2 (Table 1A, 1B).

Variety PHB18 has a greater husk extension length (4.8 cm vs 1.8 cm) than variety PHBE2 (Table 1A, 1B).

Variety PHB18 has a greater leaf length (85.2 cm vs 64.8 cm) than variety PHBE2 (Table 1A, 1B).

Variety PHB18 has a greater shank length (17.6 cm vs 10.6 cm) than variety PHBE2 (Table 1A, 1B).

Variety PHB18 has fewer primary tassel branches (8.5 vs 13.2) than variety PHBE2 (Table 1A, 1B).

Exhibit B: Novelty Statement Tables

between PHB18 and PHBE2. Each year varieties were grown in 3 locations that had different environmental conditions. Environments had Table 1A: Data from Johnston and Dallas Center, IA broken out by year and across environments are supporting evidence for differences different planting dates and were in different fields. A two-sample t-test was used to compare differences between means.

	VÁRIET	VARIETY VARIETY Count- Count-	Count- C	ount-		2	Aean S	tdDeviatS	Mean StdDeviatStdDeviat StdError- StdError- DF Po	StdError-	StdError-	DF Po	F - Value Po	Frob (2-
TRAIT	YEAR -1	-2	-		Mean-1 Mean-2	\$500 C	Diff.	ion-1	ion-2	-	2	pelo	oled	tail)_Pooled
husk length (cm)	2001PHB18	PHBE2	15	15	23.3	18.7	-4.5	1.870	1.580	0.483	0.408	28	-7.2	0.000
husk length (cm)	2002PHB18	PHBE2	15	15	23.7	19.9	-3.8	1.047	0.890	0.270	0.256	28	-10.2	0.000
husk extension length (cm)	2001PHB18	PHBE2	7	7.	4	<u>.</u>	3.1	1.957	066.0	0.505	0.256	28	5.4	0000
husk extension length					Total Control of the				***************************************					VIANATO INTERNATA LA PRINCIPATA ANTAGONIA PROPERTA PARTO DE PROPER
(cm)	2002PHB18	PHBE2	15	15	4.9	2.0	2.9	2.017	1.134	0.521	0.293	28	4.9	0.000
leaf length (cm)	2001PHB18	PHBE2	15	15	86.2	67.3	18.9	5.267	6.552	1.360	1.692	28	8.7	0.000
leaf length (cm)	2002PHB18	PHBE2	1 5	15	84.1	62.3	21.9	4.549	4.728	1.175	1.221	28	12.9	0.000
shank length (cm)	2001PHB18	PHBE2	15	15	17.1	9.7	7.3	3.731	3.173	0.963	0.819	28	5.8	0.000
shank length (cm)	2002PHB18	PHBE2	15	15	18.1	11.5	6.5	3.081	3.758	0.796	0.970	28	5.2	0.000
tassel primary branch						niae tu			********					
(# of primary		··········				en e			eria erenta atta e				•••••	
branches)	2001PHB18	PHBE2	15	15	8 .3	13.3	-5.1	1.944	3.266	0.502	0.843	28	-5.2	0.000
tassel primary branch				***			***************************************			•••••				••••
(# of primary				•••••		er com o				***************************************		•••••		
branches)	2002PHB18	PHBE2	15	15	8.7	13.1	4. 6.3	1.668	1.907	0.431	0.492	28	9.9-	0.000



Exhibit B. Novelty Statement Tables

differences between PHB18 and PHBE2. Environments had different planting dates and were in different fields. A two-sample Table 1B: Summary data from Johnston and Dallas Center, IA across years and environments are supporting evidence for t-test was used to compare differences between means.

(2- palo	0.000	0.000	0.000	0.000		0.000
Prob_(2-						***************************************
Pooled	-11.0	-7.4	14.1	7.7		-8.1
d//alie	 	58	58	8		28
E Doole	5 5 5	5	5	2		5
dError-	0.259	0.196	1.125	0.646		0.480
dError-St	0.274	0.358	0.903	0.621		0.328
/iation-St	1.418	1.073	6.163	3.538	-en-	2.631
-StdDev			0	~		(C)
StdDeviation-StdDeviation-StdError-StdError-	1.50	1.96	4.949	3.40		1.796
Aean Diff		-3.0	20.4	6.9		-4.7
<u>-</u>	19.3	1 .8	64.8	10.6		13.2
ean- Mo	23.5	4.8	85.2 (17.6		30 8.5
ount- M	30	30	30	30		30
Count-O	30	30	30	30		30
VARIETY-VARIETY-Count-Count-Mean-Mea	PHBE2	PHBE2	PHBE2	PHBE2		PHBE2
VARIETY:	PHB18	PHB18	PHB18	PHB18		PHB18
F				(ary f	nches)
TRAI	usk length (cm)	nusk extension ength (cm)	leaf length (cm)	shank length (cm)	assel primary branch (# of	primary branches) PHB18
	ੂ ਵ	로의	<u>o</u>	ठ	ट्र य	ā

EXHIBIT C:

(Corn; Maize)

(8-22-2001)

United States Department of Agriculture, Agricultural Marketing Service Science and Technology, Plant Variety Protection Office National Agricultural Library Building, Room 400 Beltsville, MD 20705-2351

> OBJECTIVE DESCRIPTION OF VARIETY CORN (Zea mays L.)

Name of Applicant(s)

Variety Name or Temporary Designation:

Pioneer Hi-Bred International, Inc

PHB18

Address (Street & No., or R.F.D. No., City,

FOR OFFICIAL USE

PVPO Number

State, Zip Code and Country)

7301 NW 62nd Avenue, P.O. Box 85, Johnston, Iowa 50131-0085

200300238

Place the appropriate number that describes the varietal characters typical of this inbred variety in the spaces below. Right justify whole numbers by adding leading zeroes if necessary. Completeness should be striven for to establish an adequate variety description. Traits designated by a "*" are considered

necessary for an adequate variety description and must be completed.

01=Light Green 02=Medium Green

COLOR CHOICES

06=Pale Yellow 07=Yellow

11=Pink 12=Light Red

16=Pale Purple 17=Purple 18=Colorless

(Use in conjunction with Munsell color code to describe all color choices; describe #25 and #26 in comments section)

21=Buff 22=Tan 23=Brown 26=Other (Describe)

03=Dark Green 04=Very Dark Green 05=Green-Yellow

08=Yellow-Orange 09=Salmon 10=Pink-Orange

13=Cherry Red 14=Red 15=Red & White

19=White 20=White Capped

24=Bronze 25=Variegated

STANDARD INBRED CHOICES (Use the most similar (in back ground and maturity) of these to make comparisons based on grow-out trial data):

Yellow Dent Families: Family

B14 B37 B73

Members CM105, A632, B64, B68 B37, B76, H84

N192, A679, B73, NC268 C103 Mo17, Va102, Va35, A682 Ch43 A619, MS71, H99, Va26 W64A, A554, A654, Pa91

Yellow Dent (Unrelated): Co109, ND246, Oh7, T232

W117, W153R W182BN White Dent

C166, H105, Ky228

Sweet Corn:

C13, Iowa 5125, P39, 2132

ı

SG1533, 4722, HP301, HP7211

Mo15W, Mo16W, Mo24W

Application Variety Data

PHB18

Standard Inbred Data **B73**

TYPE: (describe intermediate types in comments sect.)

3 (1=Sweet 2=Dent 3=Flint 4=Flour 5=Pop 6=Ornamental 7=Pipecorn)

Comments: Flint-Dent

Stand. Seed Source PI 550473

2. REGION WHERE DEVELOPED IN THE U.S.A.:

5 (1=N.West 2=N.Central 3=N.East 4=S. East 5=S. I Central, 6=S. West, 7=Other)

...... Application Variety Data **PHB18**

Page 1

Standard Variety Data

B73

(8-22-2001)

Application Variety Data PHB18		B73	Standar	d Inbre	d Data
3. MATURITY: Days Heat Units 069 1,586.0 From emergence to 50% o 67 1,541.2 From emergence to 50% o 3 92.8 From 10% to 90% pollen s From 50% silk to optimum of From 50% silk to harvest at	f plants in hed edible qu	n pollen ality	I	<u>65</u> <u>1</u>	at Units 1,486.3 1,433.3 55.5
4. PLANT:	St.Dev	S.Size	l Avg.	St.Dev	S.Size
244.6 cm Plant Height (to tassel tip)	<u>20.17</u>	<u>30</u>	l <u>233.2</u>		<u>30</u>
95.7 cm Ear Height (to base of top ear node)	<u>10.85</u>	<u>30</u>	l <u>93.6</u> I	<u>09.99</u>	<u>30</u>
16.9 cm Length of Top Ear Internode	<u>2.19</u>	<u>30</u>	l <u>15.3</u>	<u>1.93</u>	<u>30</u>
0.0 Average Number of Tillers/plant	<u>0.02</u>	30 6 6	l <u>0.0</u> l <u>1.0</u>	<u>0.02</u>	30 6 6
0.9 Average Number of Ears per Stalk	0.11	<u>6</u>	1.0	0.08	<u>6</u>
4 Anthocyanin of Brace Roots: 1=Abso	ent		<u>3</u>		
2=Faint, 3=Moderate, 4=Dark			İ		
5. LEAF:	St.Dev	S.Size	Avg. S	St.Dev	S.Size
10.0 cm Width of Ear Node Leaf	<u>1.07</u>	<u>30</u>	l <u>9.4</u>	<u>1.13</u>	30 30 30 30
85.2 cm Length of Ear Node Leaf	<u>4.95</u>	<u>30</u> <u>30</u>	82. <u>3</u>	3.03	<u>30</u>
6 Number of leaves above top ear	<u>0.65</u>	<u>30</u>	l <u>6</u>	<u>0.89</u>	<u>30</u>
19 Degrees Leaf Angle (measure from 2nd leaf above ear at anthesis to stalk above leaf)	<u>4.36</u>	<u>30</u>	l <u>18</u>	<u>4.09</u>	<u>30</u>
3 Leaf Color (Munsell code)*	5GY	34	3	7.	5GY34
2 Leaf Sheath Pubescence (Rate on sc		ᠯ:	3 3	<u></u>	1
1=none to 9=like peach fuzz)			<u> </u>		
6. TASSEL:	St.Dev	S.Size	Avg S	St.Dev	S.Size
9 Number of Primary Lateral Branches	<u>1.80</u>	<u>30</u>	<u>7.1</u>	<u>1.42</u>	<u>30</u>
32 Branch Angle from Central Spike	<u>7.90</u>	<u>30</u> <u>30</u>	<u>15.8</u>	7.27	<u>30</u> <u>30</u>
57.5 cm Tassel Length (from top leaf	<u>4.70</u>	<u>30</u>	<u>57.6</u>	<u>4.96</u>	<u>30</u>
collar node to tassel tip) 6 Pollen Shed (0=male sterile to 9=heave	av chod)		6		
1 Anther Color (Munsell code)*	2.5GY7	7 6	\ \12 \ \12	1	5Y8.54
1 Glume Color (Munsell code)*	7.5GY4	_	1	-	5GY56
1 Bar Glumes (Glume Bands): 1=Absen		 	1	-	
•			_		

Application Variety Data

Page 2

Standard Inbred Data

Note: Use chart on first page to choose color codes for color traits

**Munse11 Glossy Book of Color* (a standard color reference) Kollmorgen Inst. Corp. New Windsor NY.

(Corn; Maize)

Application Variety Data PHB18

(8-22-2001) **200300238 B73** Standard Inbred Data

Application vallety Data Phb 16		DIS	Standai	ia ilibioa	Data	
7a. EAR (Unhusked Data):	ŧ	S. Size		St.Dev S	.Size	
14 Silk Color * (3 days after emergence)	2.5R4	<u>.</u>	l <u>1 </u>	2.5GY9/4		
1 Fresh Husk Color* (25 days after 50% silking		<u> 76</u>		5GY7/8		
21 Dry Husk Color* (65 days after 50% silking	5 <u>5</u> 491	1	l <u>21</u> <u>2</u> I	2.5Y8.54		
2 Position of Ear (1=Upright, 2=Horizontal, 3	-		I <u>2</u> I <u>6</u> I 3			
5 Husk Tightness (1=very loose to 9=very tig	•		l <u>6</u>			
<u>2</u> Husk Extension: (at harvest): 1=Short (ears (<8cm), 3=Long (8-10cm to the control of the con			ا <u>غ</u>			
Long (>10cm).	.,,					
7b. EAR (Husked Ear Data):	St.Dev.	S.Size	I Avg	St.Dev S	.Size	
16.7 cm Ear Length	<u>1.47</u>	<u>30</u>	1 <u>13.3</u>	<u>1.67</u>	<u>30</u>	
44.2 mm Ear Diameter at mid-point	<u>2.47</u>	<u>30</u>	1 <u>44.8</u>	<u>2.41</u>	<u>30</u>	
137.2 gm Ear Weight	<u>29.89</u>	30 30 30 30	111.2	<u>28.55</u>	30 30 30 30	
17 Number of Kernel Rows	<u>1.34</u>	<u>30</u>	1 <u>17.6</u>	<u>1.43</u>	<u>3U</u>	
<u>2</u> Kernel Rows: 1=Indistinct 2=Distinct<u>2</u> Row Alignment:1=Straight 2=Slightly	Curved 3-	·Spiral	l <u>∠</u>			
17.6 cm Shank Length	3.40	30	l <u>≤</u> l 86	3.26	<u>30</u>	
2 Ear Taper: 1=Slight cylindrical, 2= A			<u>2</u> <u>2</u> <u>8.6</u> <u>2</u>	0.20	<u> </u>	
conical, 3=Extreme conical.	14,49,2	; ;	· <u>=</u>			
						ĺ
8. KERNEL (Dried)	St.Dev.	S.Size	l Avg. S	St.Dev. S	.Size	
8. KERNEL (Dried) 10.7 mm Kernel Length		S.Size <u>30</u>	l Avg. 8	St.Dev. S <u>0.68</u>	<u>30</u>	
	St.Dev. <u>0.78</u> <u>0.76</u>	30	1 <u>10.8</u> 1 <u>7.1</u>	St.Dev. S <u>0.68</u> <u>0.45</u>	<u>30</u>	
10.7 mm Kernel Length 8.1 mm Kernel Width 4.9 mm Kernel Thickness	0.78	30	10.8	0.68 0.45 0.53	<u>30</u>	
10.7 mm Kernel Length8.1 mm Kernel Width4.9 mm Kernel Thickness52.9 % Round Kernels (Shape Grade)	<u>0.78</u> <u>0.76</u>		1 <u>10.8</u> 1 <u>7.1</u>	<u>0.68</u> <u>0.45</u>		
 10.7 mm Kernel Length 8.1 mm Kernel Width 4.9 mm Kernel Thickness 52.9 % Round Kernels (Shape Grade) 1 Aleurone Color Pattern: 	0.78 0.76 0.52	30	1 10.8 1 7.1 3.8	0.68 0.45 0.53	<u>30</u>	
 10.7 mm Kernel Length 8.1 mm Kernel Width 4.9 mm Kernel Thickness 52.9 % Round Kernels (Shape Grade) 1 Aleurone Color Pattern: (1=Homozygous, 2=Segregating) 	0.78 0.76 0.52 18.02	30 30 30 5	1 10.8 1 7.1 3.8	0.68 0.45 0.53 8.78	30 30 30 5	1us
10.7 mm Kernel Length 8.1 mm Kernel Width 4.9 mm Kernel Thickness 52.9 % Round Kernels (Shape Grade) 1 Aleurone Color Pattern: (1=Homozygous, 2=Segregating) Aluerone Color (Munsell code)*	0.78 0.76 0.52 18.02	30 30 30 5 87/14	1 10.8 1 7.1 3.8	0.68 0.45 0.53 8.78	30 30 30 5	JMS 9110103
10.7 mm Kernel Length 8.1 mm Kernel Width 4.9 mm Kernel Thickness 52.9 % Round Kernels (Shape Grade) 1 Aleurone Color Pattern: (1=Homozygous, 2=Segregating) A Aluerone Color (Munsell code)* Hard Endosperm Color (Munsell C)*	0.78 0.76 0.52 18.02	30 30 30 5	10.8 7.1 3.8 24.5 1 .1	0.68 0.45 0.53 8.78	30 30 30 5	JMS 9/10/03
10.7 mm Kernel Length 8.1 mm Kernel Width 4.9 mm Kernel Thickness 52.9 % Round Kernels (Shape Grade) 1 Aleurone Color Pattern: (1=Homozygous, 2=Segregating) Aluerone Color (Munsell code)* Hard Endosperm Color (Munsell C)* 3 Endosperm Type:	0.78 0.76 0.52 18.02 7.5Y 7.5Y	30 30 30 5 8714 R716	1 10.8 1 7.1 3.8	0.68 0.45 0.53 8.78	30 30 30 5	JMS 9/10/03
10.7 mm Kernel Length 8.1 mm Kernel Width 4.9 mm Kernel Thickness 52.9 % Round Kernels (Shape Grade) 1 Aleurone Color Pattern: (1=Homozygous, 2=Segregating) Aluerone Color (Munsell code)* Aluerone Color (Munsell C)* 1 Endosperm Type: 1=Sweet (Su1) 2=Extra Sweet (sh2)	0.78 0.76 0.52 18.02 7.5Y 7.5Y	30 30 30 5 8714 R716	10.8 7.1 3.8 24.5 1 .1	0.68 0.45 0.53 8.78	30 30 30 5	JMS 9/10/03
10.7 mm Kernel Length 8.1 mm Kernel Width 4.9 mm Kernel Thickness 52.9 % Round Kernels (Shape Grade) 1 Aleurone Color Pattern: (1=Homozygous, 2=Segregating) Aluerone Color (Munsell code)* Hard Endosperm Color (Munsell C)* 3 Endosperm Type: 1=Sweet (Su1) 2=Extra Sweet (sh2) Starch, 4=High Amylose Starch, 5=W	0.78 0.76 0.52 18.02 7.5Y 7.5Y 3=Normal axy Starch	30 30 30 5 5 R714 R716	10.8 7.1 3.8 24.5 1 .1	0.68 0.45 0.53 8.78	30 30 30 5	JMS 9/10/03
10.7 mm Kernel Length 8.1 mm Kernel Width 4.9 mm Kernel Thickness 52.9 % Round Kernels (Shape Grade) 1 Aleurone Color Pattern: (1=Homozygous, 2=Segregating) Aluerone Color (Munsell code)* Aluerone Color (Munsell C)* 1 Endosperm Type: 1=Sweet (Su1) 2=Extra Sweet (sh2)	0.78 0.76 0.52 18.02 7.5Y 7.5Y 3=Normal axy Starch	30 30 30 5 5 R714 R716	10.8 7.1 3.8 24.5 1 .1	0.68 0.45 0.53 8.78	30 30 30 5	JMS 9/10/03
10.7 mm Kernel Length 8.1 mm Kernel Width 4.9 mm Kernel Thickness 52.9 % Round Kernels (Shape Grade) 1 Aleurone Color Pattern: (1=Homozygous, 2=Segregating) Aluerone Color (Munsell code)* Aluerone Color (Munsell C)* 1 Endosperm Type: 1=Sweet (Su1) 2=Extra Sweet (sh2) Starch, 4=High Amylose Starch, 5=W 6=High Protein, 7=High Lysine, 8=Su	0.78 0.76 0.52 18.02 7.5Y 7.5Y 3=Normal axy Starch	30 30 30 5 5 R714 R716	10.8 7.1 3.8 24.5 1 .1	0.68 0.45 0.53 8.78	30 30 30 5	JMS 9/10/03
10.7 mm Kernel Length 8.1 mm Kernel Width 4.9 mm Kernel Thickness 52.9 % Round Kernels (Shape Grade) 1 Aleurone Color Pattern: (1=Homozygous, 2=Segregating) Aluerone Color (Munsell code)* Hard Endosperm Color (Munsell C)* 3 Endosperm Type: 1=Sweet (Su1) 2=Extra Sweet (sh2) Starch, 4=High Amylose Starch, 5=W 6=High Protein, 7=High Lysine, 8=Su 9=High Oil, 10=Other	0.78 0.76 0.52 18.02 7.5Y 7.5Y 3=Normal axy Starch per Sweet	30 30 30 5 87/14 R7/16	10.8 7.1 3.8 24.5 1 27.2 7 3	0.68 0.45 0.53 8.78 2.5Y8 10YR7	30 30 30 5 2 12	JMS 9/10/03
10.7 mm Kernel Length 8.1 mm Kernel Width 4.9 mm Kernel Thickness 52.9 % Round Kernels (Shape Grade) 1 Aleurone Color Pattern: (1=Homozygous, 2=Segregating) Aluerone Color (Munsell code)* Hard Endosperm Color (Munsell C)* 3 Endosperm Type: 1=Sweet (Su1) 2=Extra Sweet (sh2) Starch, 4=High Amylose Starch, 5=W 6=High Protein, 7=High Lysine, 8=Su 9=High Oil, 10=Other 29.0 gm Weight per 100 Kernels	0.78 0.76 0.52 18.02 7.5Y 7.5Y 3=Normal axy Starch per Sweet	30 30 30 5 87/14 R7/16	10.8 7.1 3.8 24.5 1 27.2 7 3	0.68 0.45 0.53 8.78 2.5Y8 10YR7	30 30 30 5 2 12	JMS 9/10/03
10.7 mm Kernel Length 8.1 mm Kernel Width 4.9 mm Kernel Thickness 52.9 % Round Kernels (Shape Grade) 1 Aleurone Color Pattern: (1=Homozygous, 2=Segregating) A Aluerone Color (Munsell code)* Hard Endosperm Color (Munsell C)* 3 Endosperm Type: 1=Sweet (Su1) 2=Extra Sweet (sh2) Starch, 4=High Amylose Starch, 5=W 6=High Protein, 7=High Lysine, 8=Su 9=High Oil, 10=Other 29.0 gm Weight per 100 Kernels (Unsized sample) 9. COB: 27.5 mm Cob Diameter at mid-point	0.78 0.76 0.52 18.02 7.5Y 7.5Y 3=Normal axy Starch per Sweet 2.92	30 30 5 5 R714 R716 (se),	10.8 7.1 3.8 24.5 1 27.2 7 3	0.68 0.45 0.53 8.78 2.5Y8 10YR7	30 30 30 5 2 12	JMS 9/10/03
10.7 mm Kernel Length 8.1 mm Kernel Width 4.9 mm Kernel Thickness 52.9 % Round Kernels (Shape Grade) 1 Aleurone Color Pattern: (1=Homozygous, 2=Segregating) Aluerone Color (Munsell code)* Hard Endosperm Color (Munsell C)* 3 Endosperm Type: 1=Sweet (Su1) 2=Extra Sweet (sh2) Starch, 4=High Amylose Starch, 5=W 6=High Protein, 7=High Lysine, 8=Su 9=High Oil, 10=Other 29.0 gm Weight per 100 Kernels (Unsized sample)	0.78 0.76 0.52 18.02 7.5Y 7.5Y 3=Normal axy Starch per Sweet 2.92	30 30 5 5 R714 R716 (se),	10.8 7.1 3.8 24.5 .1 07.8 7 3	0.68 0.45 0.53 8.78 2.5Y8 10YR7	30 30 30 5 2 12 5	JMS 9/10/03

EXHIBIT C:

(Corn; Maize) (8-22-2001) Standard Inbred Data

Application Variety Data PHB18 B73 10. DISEASE RESISTANCE: (1-most susceptable to 9-most resistant, Leave blank if not tested, Leave race or strain options blank if polygenic) A. LEAF BLIGHTS, WILTS, AND LOCAL INFECTION DISEASES Anthracnose Leaf Blight (Colletotrichum graminicola) Common Rust (Puccinia sorghi) Common Smut (Ustilago maydis) Eyespot (Kabatiella zeae) Gross's Wilt (Clavibacter michiganense spp. nebraskense) 4 Gray Leaf Spot (Cercospora zeae-maydis) <u>3</u> Helminthosporium Leaf Spot (Bipolaris zeicola) Race: <u>3</u> 6 Northern Leaf Blight (Exserohilum turcicum) Race: Southern Leaf Blight (Bipolaris maydis) Race: <u>2</u> 2 4 Southern Rust (Puccinia polysora) 5 Stewart's Wilt (Erwinia stewartii) Other (Specify): **B. SYSTEMIC DISEASES** Corn Lethal Necrosis (MCMV and MDMV) Head Smut (Sphacelotheca reiliana) Maize Chlorotic Dwarf Virus (MDV) _ Maize Chlorotic Mottle Virus (MCMV) Maize Dwarf Mosaic Virus (MDMV) Sorghum Downy Mildew of Corn (Peronosclerospora sorghi) Other (Specify): C. STALK ROTS 5 Anthracnose Stalk Rot (Colletotrichum graminicola) __ Diploidia Stalk Rot (Stenocarpella mavdis) __ Fusarium Stalk Rot (Fusarium moniliforme) __ Gibberella Stalk Rot (Gibberella zeae) Other (Specify): D. EAR AND KERNEL ROTS Aspergillus Ear and Kernel Rot (Aspergillus flavus) 8 Diplodia Ear Rot (Stenocarpella maydis) 4 Fusarium Ear and Kernel Rot (Fusarium moniliforme) Gibberella Ear Rot (Gibberella zeae) Other (Specify): tion Variety Data Page 4 Standard Inbred Data Application Variety Data **PHB18 B73**

Standard Inbred Data

Application Variety Data

EXHIBIT C:

PHB18 11. INSECT RESISTANCE: (1-most susceptable to 9-most resistant) ı __ Banks Grass Mite (Oligonychus pratensis) __ Corn Earworm (Helicoverpa zea) Leaf Feeding __ Silk Feeding ___ . _mg larval wt. __ Ear Damage __ Corn Leaf Aphid (Rhopalosiphum maidis) Corn Sap Beetle (Capophilus dimidiatus) _ European Corn Borer (Ostrinia nubilalis) 1st. Generation (Typically whorl leaf feeding) 2nd. Generation (Typically leaf sheath-collar feeding) __ Stalk Tunneling ____ . _cm tunneled/plant _ Fall Armyworm (Spodoptera frugiperda) _ Leaf Feeding _ Silk Feeding: ___, mg larval wt. _ Maize Weevil (Sitophilus zea maize) __ Northern Rootworm (Diabrotica barberi) _ Southern Rootworm (Diabrotica undecimpunctata) _ Southwestern Corn Borer (Diatreaea grandiosella) _ Leaf Feeding __ Stalk Tunneling ____. _cm tunneled/plant __ Two-spotted Spider Mite (Tetranychus utricae) _ Western Rootworm (Diabrotica virgifera virgifera) _ Other (Specity) 12. AGRONOMIC TRAITS: 5 Staygreen (at 65 days after anthesis; rate from 1-worst to excellent) % Dropped Ears (at 65 days after anthesis) % Pre-anthesis Brittle Snapping 22 % Pre-anthesis Root Lodging 2 % Post-anthesis Root Lodging (at 65 days after anthesis) % Post-anthesis Stalk Lodging 4,805.0 Kg/ha Yield of inbred per se (at 12-13% grain moisture) 5,177.0 13 MOLECULAR MARKERS: (0-data unavailable; 1-data available but not supplied; 2-data supplied) Isozymes 1 RFLP's RAPD's Application Variety Data Page 5 Standard Inbred Data **PHB18 B73**

COMMENTS: (eg. state how heat units were calculated, standard inbred seed source, and/or where data was collected. Continue in exhibit D)

Please note the data presented in Exhibit B and C, "Objective Description of Variety," are collected primarily at Johnston and Dallas Center, Iowa. The data in Tables 1A and 1B are from two sample t-tests using data collected in Johnston and Dallas Center, IA. These traits in exhibit B collectively show distinct differences between the two varieties.

The data collected in exhibit C was collected in 2001 and 2002 for page 1 and 2. There were 3 different planting dates planted for these trials. There are environmental factors that differ from year to year and from planting date to planting date. Environmental temperature and precipitation differences during the vegetative and grain fill periods can impact plant and grain traits, and are a source of variability. The environmental conditions described above could result in larger standard deviations. The variation associated with environment to environment is normally higher than the variation associated within locations. Also, the ear and sizing traits can vary depending on how well pollinated the ears are and how consistent the weather is during the grain fill period. I have enclosed a table that shows monthly temperature and precipitation in 2001 and 2002.

Exhibit 1. Temperature and Precipitation differences from Ankeny, IA

TEMPERATURE

YEAR	MAY	JUN	JULY	AUG	AVERAGE
1994	59.8	70.7	71.9	69.0	67.9
1995	56.2	69.4	74.3	76.9	69.2
1996	56.2	69.3	71.3	70.5	66.8
1997	53.5	70.6	74.1	69.6	67.0
1998	64.7	66.6	74.8	73.5	69.9
1999	60.7	69.7	78.7	70.5	69.9
2000	63.5	68.9	73.2	74.2	70.0
2001	61.3	69.0	76.7	74.2	70.3
2002	57.7	73.5	77.9	71.7	70.2

RAINFALL

YEAR	MAY	JUN	JULY	AUG	Total
1994	3.67	5.75	1.71	4.18	15.31
1995	5.04	4.19	2.94	2.87	15.04
1996	8.47	4.35	2.51	2.14	17.47
1997	4.32	3.27	4.10	1.36	13.05
1998	6.46	11.07	5.70	4.96	28.19
1999	6.46	4.54	4.45	6.55	21.85
2000	5.40	5.80	3.16	1.78	16.14
2001	5.72	3.87	2.05	1.92	13.56
2002	2.91	2.78	5.34	4.00	15.03

U.S. DEPARTMENT OF AGRICULTURE AGRICULTURAL MARKETING SERVICE	The following statements are made in account 1974 (5 U. S. C. 552a) and the Paperwork				
EXHIBIT E STATEMENT OF THE BASIS OF OWNERSHIP	Application is required in order to determ certificate is to be issued (7 U.S.C. 2421). until certificate is issued (7 U.S.C. 2426).				
1.NAME OF APPLICANT(S)	2.TEMPORARY DESIGNATION OR EXPERIMENTAL NUMBER	3. VARIETY NAME			
PIONEER HI-BRED INTERNATIONAL, INC.	OR EXPERIMENTAL NOWIDER	PHB18			
4 .ADDRESS (Street and No., or R.F.D. No., City, State, and ZIP, and Country)	5.TELEPHONE (include area code)	6. FAX (include area code)			
7301 NW 62 nd AVENUE P.O.BOX 85	515-270-4051	515-253-2125			
JOHNSTON, IA 50131-0085 200300238					
8. Does the applicant own all rights to the variety? Mark an "X" in appropriate block	k. If no, please explain: 🛛 YES 🗆	NO			
9.1s the applicant (individual or company) a U.S. national or U.S. based company?	P ⊠ YES □ NO				
If no, give name of country					
10. Is the applicant the original owner? YES NO If no, please ar	nswer <u>one</u> of the following:				
a. If original rights to variety were owned by individual(s), is(are) the original	al owner(s) a U.S. national(s)?				
☐ YES ☐ NO if no, give name of country					
b. If original rights to variety were owned by a company(ies), is(are) the original	ginal owner(s) a U.S. based company?				
	gillar emisi(e) a ere, saces esimpany				
23 120 El No Il no, give name of country					
11. Additional explanation on ownership (if needed, use reverse for extra space):					
PHB18 is owned by Pioneer Hi-Bred International, Inc.					
Pioneer Hi-Bred International, Inc. (PHI), Des Moines, Iowa, and/or its whol Moines, Iowa, is the employer of the plant breeders involved in the selection Pioneer Overseas Corporation has the sole rights and ownership of PHB18 p and/or POC at the time such variety was created. Individuals do not retain an	and development of PHB18. Pioneer Hi-Bred ursuant to written contracts that assign all rights	International and/or			
PLEASE NOTE:					
Plant variety protection can be afforded only to owners (not licensees) who meet one of the	following criteria:				
1. If the rights to the variety are owned by the original breeder, that person must be a U. Which affords similar protection to nationals of the U.S. for the same genus and spec		ry, or national of a country			
2. If the rights to the variety are owned by the company which employed the original bre country, or owned by national of a country which affords similar protection to national		ed by nationals of a UPOV member			
3. If the applicant is an owner who is not the original owner, both the original owner and	d the applicant must meet one of the above crite	жіа.			
The original breeder/owner may be the individual or company who directed final breeding.	See section 41(a)(2) of the Plant Variety Prote	ection Act for definition.			
According to the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information collection is 0581-0055. The time required to compete this information collection is estima existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection.	ated to average 10 minutes per response, including the				

The U.S. Department of Agriculture (USDA) prohibits discrimination in its programs on the basis of race, color, national origin, sex, religion, age, disability, political beliefs, and marital or familial status (Not all prohibited bases apply to all programs). Persons with disabilities who require alternative means for communication of program information (braille, large print, audiotape, etc.) should contact USDA's TARGET Center at 202-720-2600 (voice and TDD).

To file a complaint, write Secretary of Agriculture, U.S. Department of Agriculture, Washington, D.C. 20250, or call 1-800-245-6340 (voice) or (202) 720-1127 (TDD) USDA is an equal employment

STD-470-E (07-97) (Destroy previous editions).
Electronic version designed using WordPerfect In Forms by USDA-AMS-IMB

opportunity employer.